

330732 to 330921—Continued

330811. Col. No. P-519. From Tinfa in Nekempt. Local name Anhero. Collected by Agricultural Agent in area. Used for injera. Seed pearly white. Head semicompact, no central rachis. Glumes black, cover $\frac{1}{2}$ of seed.
330812. Col. No. P-520. From Nekempt area. Local name Bobay. Collected by agricultural agent in area. Used for porridge, injera, and tella. Seed white. Head long, semicompact, central rachis. Glumes pubescent, covering $\frac{2}{3}$ of seed, awned.
330813. Col. No. P-521. From Sire, Nekempt. Local name Bobay. Collected by agricultural agent in area. Used for porridge, injera, and tella. Seed white. Head long, semicompact, central rachis. Glumes pubescent, covering $\frac{2}{3}$ of seed, awned.
330814. Col. No. P-522. From Sire, Nekempt. Collected by agricultural agent. Seed yellow. Head very compact, durra type. Glumes cover $\frac{1}{2}$ of seed, awned.
- From Nekempt. Local name Bisinga. Collected by agricultural agent.
330815. Col. No. P-523. Seed reddish brown. Head open, reduced central rachis. Glumes cover $\frac{1}{2}$ of seed, straw to brown, awned.
330816. Col. No. P-524. Seed deep red. Head open, no central rachis. Glumes cover $\frac{1}{2}$ of seed, dark, awns.
330817. Col. No. P-525. Seed grey brown. Head open, no central axis. Glumes cover $\frac{1}{2}$ of seed, dark, awned. Used for injera and tella.
330818. Col. No. P-526. From Tinfa, Nekempt. Local name Dappo. Collected by agricultural agent. Used for tella and injera. Seed reddish. Head semicompact, no central rachis. Glumes cover $\frac{1}{2}$ of seed, mahogany, awnless.
330819. Col. No. P-527. From Nekempt area. Local name Sippo. Collected by agricultural agent. Used for tella and porridge. Seed brown. Head open, central rachis. Glumes covering seed, black, awnless.
330820. Col. No. P-528. From Arjo. Local name Red Bisinga. Collected by agricultural agent. Seed reddish brown. Head semicompact, no central rachis. Glumes cover $\frac{1}{2}$ of seed, red to brown, awns.
330821. Col. No. P-529. From Nekempt area. Local name Bisinga. Collected by agricultural agent. Seed reddish brown. Heads semicompact, reduced rachis. Glumes cover $\frac{1}{2}$ of seed, awned.
330822. Col. No. P-530. From Masara, Nekempt. Local name Chabbi. Collected by agricultural agent. Used for injera. Seed white. Heads long, open, central rachis, extended. Glumes cover $\frac{2}{3}$ of seed, awned.
330823. Col. No. P-531. From Nekempt area. Local name Sippo. Collected by agricultural agent. Used for porridge and tella. Seed brown. Heads open, sudangrass type. Glumes covering seed, purple black, awned.
330824. Col. No. P-532. Near the Didessa River, Didessa, Nekempt. Local name Red Dappo. Collected by agricultural agent. Used for porridge, injera, and tella. Seed reddish brown. Head open, to 20 cm. long pedicels, seed clustered on end. Glumes cover $\frac{1}{2}$ of seed, black, pubescent, awnless.
330825. Col. No. P-533. From Nekempt area. Collected by agricultural agent. Seed reddish brown. Heads open, central rachis. Glumes cover $\frac{1}{2}$ of seed, brown, awned.
330826. Col. No. P-534. From Arjo area. Local name Boba. Collected by agricultural agent. Seed white. Heads open, no rachis. Glumes cover $\frac{1}{2}$ of seed, sienna, awnless.