

**383326 to 383338—Continued**

entrance gate. Elevation 1,500 m. Wet drainage line in vlei. Strongly creeping perennial. Culms compressed, each with several flowering branches, up to 130 cm. long. Leaves with blades expanded at first, folding very readily, blunt at the apex. Inflorescence a solitary raceme, spikelike, compressed, awnless. Black vlei clay soil. Common in large patches. January 1968.

**383332.** *HYLEBATES CORDATUS* Chippind. Gramineae.

848. Townlands on main road to Beira, near Eastlands farm boundary. Elevation 1,000 m. On old anthill in mixed Msasa woodland. Heavy clay and black leaf-mold soil. February 1969.

**383333.** *MISCANTHIDIUM VIOLACEUM* (K. Schum.) Robyns Gramineae.

957. West of turnoff to farm De Rots. Elevation 1,554 m. Streambank in center of vlei. Black clay soil. Common. January 1971.

**383334.** *OLYRA LATIFOLIA* L. Gramineae.

1088. Northeast edge of Chirunda Forest. Elevation 1,000 m. Closed canopy indigenous forest, a tangle of undergrowth at edge of break in the forest canopy where *Khya nyassica* is common. Forest soils derived from dolerite. Very common in local patch under *Khya*. May 1973.

**383335.** *PANICUM DREGEANUM* Nees Gramineae.

Rusape Townlands near electric transformer station. Elevation 1,400 m. Very shallow, poor granite sandveld, partially waterlogged soil. Locally occasional. November 1972.

**383336.** *RHYNCHELYTRUM RHODESIANUM* (Rendle) Stapf & C. E. Hubb. Gramineae.

1058. Rusape Townlands near electric transformer station. Elevation 1,400 m. Densely tufted perennial. Culms 50 to 90 cm. high, erect, nodes conspicuously bearded. Leaf blades up to 2 mm. wide, rolled at first, later expanded. Inflorescence a narrow panicle. Spikelets with long hairs mainly on upper part. Disturbed veld between main road and electric transformer station. Very shallow, poor, granite sandveld soil. October 1972.

**383337.** *SPOROBOLUS PANICOIDES* A. Rich. Gramineae. **Dropseed.**

Granite whaleback (ruwari) Chitsanga, near Rowa School. Elevation 1,200 m. Shallow soils fringing the woodland near top of exposed granite. Detritus and shallow sand soil. Locally common. March 1973.

**383338.** *TRACHYPOGON SPICATUS* (L. f.) Ktze. Compositae.

849. Umtali Heights. Elevation 1,600 m. Perennial, densely tufted, mostly 70 to 100 cm. high, erect. Leaf blades 2 mm. wide, tapering to a hair point. Inflorescence usually one, rarely two spikelike racemes, dense and hairy, 10 to 17 cm. long, with conspicuous, geniculate, hairy awns. Open highveld grassland. Brown clays from serpentine soil. Locally common. February 1969.

**383339 to 383341.**

From Iran. Seed collected by J. L. Schwendiman, Plant Materials Center, Soil Conservation Service, Pullman, Washington. Received September 15, 1972.

**383339.** *VICIA ERVILIA* (L.) Willd. Leguminosae. **Bitter vetch.**

351. Nujian. Annual to 25 cm. tall. Constricted pods. A heavy seeder, does not shatter. Seeded on watershed area. Produced a dense stand and a good seed crop. Wild.

**383340 and 383341.** *VICIA TENUIFOLIA* Roth