

PI 543212. *Citrullus lanatus* (Thunb.) Matsum. & Nakai CUCURBITACEAE
Watermelon

Donated by: Williams, D.E., New York Botanical Garden, Bronx, New York, United States. **remarks:** Collected in Northern Bolivia, September 8, 1988 through December 12, 1988. Received November 1989.

donor id: 807. **origin:** Bolivia. **local name:** Sandia.
collected: November 08, 1988. **collector id:** 807.
locality: 21 de Septiembre, small village on L bank of Rio Beni, Pando Department, Madre de Dios Province.
latitude: 11 deg. 25 min. S. **longitude:** 66 deg. 30 min. W. **elevation:** 135m. **remarks:** Teardrop-shaped fruit 40cm long, 20cm diameter, green with lighter green spots in longitudinal lines. Locally cultivated watermelon. Cultivated. Seed.

PI 543213 to 543214. *Cyclanthera pedata* (L.) Schrad. CUCURBITACEAE
Achocha

Donated by: Williams, D.E., New York Botanical Garden, Bronx, New York, United States. **remarks:** Collected in Northern Bolivia, September 8, 1988 through December 12, 1988. Received November 1989.

PI 543213 **donor id:** 826. **origin:** Bolivia. **local name:** Caiba, Maxixi peruano, Achoscha (all Spanish). **collected:** November 13, 1988. **collector id:** 826. **locality:** Cobija, market, Pando Department, Nicolas Suarez Province.
latitude: 11 deg. 00 min. S. **longitude:** 68 deg. 45 min. W. **elevation:** 200m. **remarks:** Fruit oblong, pointed at both ends, somewhat flattened, smooth, waxy, partially ribbed, to 14cm long, 6cm wide, hollow, light green turning yellowish. Seeds black, roughly textured, somewhat triangular, dentate, about 10 borne on an axially free placenta. Cultivated locally as a vegetable for soups, stews, and stuffing. Cultivated. Seed.

PI 543214 **donor id:** 872. **origin:** Bolivia. **local name:** Achoscha. **collected:** November 27, 1988. **collector id:** 872. **locality:** Pompeya market, Trinidad, Beni Department, Cercado Province. **latitude:** 14 deg. 48 min. S. **longitude:** 64 deg. 52 min. W. **elevation:** 155m. **remarks:** Fruit oblong, to 18cm long, 6cm wide, acuminate, flattened, shallowly ribbed, soft, hollow, light green. Seeds black, vaguely triangular, dentate, rough-textured, 8-14 borne on an axially free placenta. Cucumber-like odor. Cultivated as a vegetable for soups and for stuffing. Cultivated. Seed.