

The following were developed by R. Veitch, Bureau of Plant Research, Department of Agriculture, Brisbane, Queensland, Australia. Donated by A.I. Campbell, Long Ashton Research Station, Long Ashton, Bristol, England B518 9AF, United Kingdom. Received 03/31/1987.

PI 589700. *Malus domestica* Borkh.

Rev. W. Wilks. Pedigree - Peasgood's Nonsuch and Ribston Pippin; recorded 1904. introduced 1908. Comments:: Size large 88:76 mm; shape flat, conic, convex, slightly ribbed at eye and on body; skin pale yellow with slight flush and stripes; flesh, cooks well, tender, white; flavor acid; season mid. Tree: rather small spreading. Produces spurs very freely. Very handsome, mid-season cooking apple. Primarily grown for garden and exhibition. Tends to be biennial.

The following were developed by T. Squire. Donated by A.I. Campbell, Long Ashton Research Station, Long Ashton, Bristol, England B518 9AF, United Kingdom. Received 03/31/1987.

PI 589701. *Malus domestica* Borkh.

Lane's Prince Albert. Pedigree - Said to be Russet Nonpariel x Dumelow's Seedling. Raised around 1840; introduced by Lane around 1850. Comments:: Size large 76:59 mm; shape intermediate, truncate-conic, convex, slightly ribbed at eye, skin yellowish green, flushed orange, striped red; flesh very soft, fine, white tinged green; flavor acid; season very late. Tree: very vigorous. Well known, late keeping apple. Primarily a garden variety, but also grown on a small scale commercially in UK. A regular cropper of compact habit.

The following were donated by A.I. Campbell, Long Ashton Research Station, Long Ashton, Bristol, England B518 9AF, United Kingdom. Received 03/31/1987.

PI 589702. *Malus domestica* Borkh.

Early Crimson.

PI 589703. *Malus domestica* Borkh.

Kingston Black. Comments:: Famous English cider apple - one of only three English varieties used as a single-variety cider. Classed as bitter sharp. Medium size, round fruit. Crimson to purplish over yellowish orange background color. Strongly astringent after-taste. Makes a distinctively flavored cider without blending. Vigorous rounded tree. Somewhat hard to grow. Ripens from mid-September to mid-October depending on location. Subject to scab and canker.

The following were developed by J. Musch. Donated by G. Gilles, Research Station of Gorsem, Virology Dept., Brede Akker3, B3800 Sint Truiden, Belgium. Received 03/18/1987.

PI 589704. *Malus domestica* Borkh.

Joseph Musch. Pedigree - In NFT collection is identical with Gascoyne's Scarlet. Recorded 1872. Comments:: Size large; shape variable, intermediate, truncate-conic, convex, ribbed at eye and on body; skin pale yellow with bright red flush and stripes, russet at base; flesh rather coarse, dry, whitish yellow; flavor subacid, sweet. Season mid to late.

The following were donated by P.J. Sampson, Dept. of Agriculture Tasmania, Senior Plant Pathology Section, 1 Franklin Wharf, Hobart, Tasmania, Australia. Received 05/22/1987.