

The following were donated by Irrigated Agric. Research Ext. Center, P.O. Box 30, Washington State University, Prosser, Washington 99350, United States. Received 02/15/1988.

PI 589899. *Malus domestica* Borkh.
Red Baron.

The following were developed by Dummer. Donated by Irrigated Agric. Research Ext. Center, P.O. Box 30, Washington State University, Prosser, Washington 99350, United States. Received 02/15/1988.

PI 589900. *Malus domestica* Borkh.
Discovery. Collected in Unknown. Originated in England around 1900. Pedigree - Worcester Pearmain x ? Beauty of Bath; Commercial Grower No. 2345 p. 725 IRHS; Raised circa 1949; transplanted 1955. Pollinate with Gravenstein Lodi or Summer Red. Comments:: Size medium to large 60-77:45-61 mm; shape flat, rectangular to truncate-conic, convex, slight ribbing at eye, very even; skin pale greenish yellow, flushed bright red, light dots, fine scattered russet, solid russet at base. Flesh crisp, yellowish white tinged pink, excellent flavor. Some resistance to scab. Ripens from early to mid-August depending on location. Not a keeper. fruit tends to crack. Hardy compact tree - natural semi-dwarf. Flowers frost resistant. Additional LIT.CIT. 1992-93. H. Apple. Tsolum River Fruit Trees Catalogue. p. 16.

The following were developed by Univ. of Minnesota Fruit Breeding Farm, Excelsior, Minnesota, United States. Donated by Irrigated Agric. Research Ext. Center, P.O. Box 30, Washington State University, Prosser, Washington 99350, United States. Received 02/15/1988.

PI 589901. *Malus domestica* Borkh.
Redwell. Pedigree - Scott's Winter open-pollinated. Seed sown circa 1911, selected 1923, introduced 1946; tested as Minn. 638. Comments:: Size medium to large 67-76;:57-69 mm; shape tall, sometimes flat, rectangular, sometimes truncate-conic, convex, sometimes ribbed at eye and slightly on body; skin greenish yellow, flushed reddish orange, carmine streaks, russet at base, russet dots, patches and netting; flesh a little soft, greenish white; flavor subacid, mild, excellent; season mid-late. Vigorous tree - strong framework, bears annually. Partially dwarfing; medium susceptibility to cedar rust and scab. Storage until Jan.

The following were donated by Irrigated Agric. Research Ext. Center, P.O. Box 30, Washington State University, Prosser, Washington 99350, United States. Received 02/15/1988.

PI 589902. *Malus domestica* Borkh.
Regent.

PI 589903. *Malus domestica* Borkh.
Smokehouse. Collected in Unknown. Originated on the Farm of William Gibbons, Lancaster County, near Millcreek, PA. Pedigree - Before 1837. Thought to be a seedling of Vandevere of Delaware and Pennsylvania. Comments:: Size large 84;64 mm; shape flat, rectangular to truncate-conic, convex, not ribbed; skin yellow or greenish, mottled dull red, sometimes flushed bright red, striped carmine, russet dots; flesh rather firm, moderately fine and tender, crisp, slightly tinged yellow, flavor subacid, slightly aromatic; season mid to very late. Vigorous tree, requires pruning. Reliable. Subject to apple scab. Fresh cider