

**PI 590028. Malus halliana Koehne**

Collected 1980 in China. Shennongia Forest District, northwestern part of Hubei Province bounded on the west by Sichuan Province and on the South by the confluence of the Juichong and Dangyang Rivers. Pedigree - Wild. Comments:: Participants of the 1980 Sino-American Botanical Expedition observed a varied climate ranging from warm temperate to subtropical. A complex topography of steep mountainous terrain supports rich and varied flora (including relict families, genera and species).

**PI 590029. Malus halliana Koehne**

Collected 1980 in China. Shennongia Forest District, northwestern part of Hubei Province, bounded on the west by Sichuan Province and on the south by the confluence of the Juichong and Dangyang Rivers. Pedigree - Wild. Comments:: Participants of the 1980 Sino-American Botanical Expedition observed a varied climate ranging from warm temperate to subtropical. A complex topography of steep mountainous terrain supports rich and varied flora (including relict families, genera and species).

**PI 590030. Malus halliana Koehne**

Collected 1980 in China. Shennongia Forest District, northwestern part of Hubei Province, bounded on the west by Sichuan Province and on the south by the confluence of the Juichong and Dangyang Rivers. Pedigree - Wild. Comments:: Participants of the 1980 Sino-American Botanical Expedition observed a varied climate ranging from warm temperate to subtropical. A complex topography of steep mountainous terrain supports rich and varied flora (including relict families, genera and species).

**PI 590031. Malus halliana Koehne**

Collected 1980 in China. Shennongia Forest District, northwestern part of Hubei Province, bounded on the west by Sichuan Province and on the south by the confluence of the Juichong and Dangyang Rivers. Pedigree - Wild. Comments:: Participants of the 1980 Sino-American Botanical Expedition observed a varied climate ranging from warm temperate to subtropical. A complex topography of steep mountainous terrain supports rich and varied flora (including relict families, genera and species).

**PI 590032. Malus halliana Koehne**

Collected 1980 in China. Shennongia Forest District, northwestern part of Hubei Province, bounded on the west by Sichuan Province and on the south by the confluence of the Juichong and Dangyang Rivers. Pedigree - Wild. Comments:: Participants of the 1980 Sino-American Botanical Expedition observed a varied climate ranging from warm temperate to subtropical. A complex topography of steep mountainous terrain supports rich and varied flora (including relict families, genera, and species).

The following were collected by H.J. Kelsey. Donated by Rob Nicholson, The Arnold Arboretum of Harvard Univ., The Arborway, Jamaica Plain, Massachusetts 02130, United States. Received 03/22/1989.

**PI 590033. Malus floribunda Siebold ex Van Houtte**

Collected 1927 in Unknown. Pedigree - Wild.

**PI 590034. Malus floribunda Siebold ex Van Houtte**

Collected 1927 in Unknown. Pedigree - Wild.

The following were collected by Karl Sax. Donated by Rob Nicholson, The Arnold Arboretum of Harvard Univ., The Arborway, Jamaica Plain, Massachusetts 02130, United States. Received 03/22/1989.

**PI 590035. Malus floribunda Siebold ex Van Houtte**

Collected 1953 in Unknown. Pedigree - Wild.