

1680 Madison, Wooster, Ohio 44691-4096, United States; S.K. St. Martin, Ohio Agr. Exp. Sta., Ohio State University, Department of Agronomy, Wooster, Ohio 44691-4096, United States; A.J. Calip-DuBois, Ohio State University, Ohio Agricultural Res. and Develop. Ctr., Dept. of Horticulture and Crop Science, Columbus, Ohio 43210, United States; Ron Fioritto, Ohio State University, Ohio Agricultural Research & Development Center, Dept. of Horticulture & Crop Science, Wooster, Ohio 44691, United States; A.F. Schmitthenner, Ohio State University, Dept. of Plant Pathology, Columbus, Ohio 43210, United States; R.J. Martin, USDA, ARS, Ohio State University, Dept. of Food Science and Technology, Columbus, Ohio 43210, United States. Received 09/30/1996.

PI 595843. *Glycine max* (L.) Merr.

Cultivar. Pureline. "Flint". CV-362; PVP 9600408. Pedigree - GR8836 x Elgin 87. Maturity group II, growth indeterminate. Flowers purple, pubescence tawny, pods brown. Seeds dull yellow, hila black, mean content of 41.1% protein and 20.6% oil, size 16 mg seed-1. Excellent tolerance to *Phytophthora* root rot and carries Rps1-k allele which confers multi race resistance to *Phytophthora sojae*. Moderately susceptible to brown stem rot and susceptible to soybean cyst nematode.

The following were developed by Holden's Foundation Seeds, Inc, United States . Received 10/02/1996.

PI 595844. *Zea mays* L. ssp. *mays*

Cultivar. "LH179". PVP 9700002.

PI 595845. *Zea mays* L. ssp. *mays*

Cultivar. "LH236". PVP 9700003.

PI 595846. *Zea mays* L. ssp. *mays*

Cultivar. "LH281". PVP 9700004.

PI 595847. *Zea mays* L. ssp. *mays*

Cultivar. "LH290". PVP 9700005.

The following were developed by Mark E. Sorrells, Cornell University, Dept. of Plant Breeding & Biometry, 252 Emerson Hall, Ithaca, New York 14853-1902, United States. Received 10/10/1996.

PI 595848. *Triticum aestivum* L., nom. cons.

Cultivar. Pureline. "CAYUGA"; NY262-37-422. PVP 9700007. Pedigree - Geneva/Clark's Cream//Geneva. Soft white winter wheat. At booting flag leaf erect and twisted, stems waxy bloom and yellow-green, leaves yellow-green. At maturity stems hollow white. Spikes mid-dense, tapering, apically awnletted, length avg. 7-8 cm. Glumes white, length medium, wide, shoulder rounded, beak acute. Kernels ovate, cheeks rounded, brush medium, crease shallow and mid-deep. Seed length avg 6.3 mm, width 3.6 mm, 35 g/1000 seed. Resistance to preharvest sprouting. Moderate resistance to loose smut, powdery mildew, and wheat spindle streak mosaic virus. Susceptible to leaf rust and stripe rust.

The following were developed by C.F. Konzak, Washington State University,