

The following were developed by Dennis Thomas, University of Illinois, Department of Agronomy, 1102 S. Goodwin Avenue, Urbana, Illinois 61801, United States; Cecil D. Nickell, University of Illinois, Department of Crop Science, 1102 S. Goodwin Av., Urbana, Illinois 61801, United States; T.R. Cary, University of Illinois, Illinois Agr. Exp. Sta., Dept. of Agronomy, Urbana, Illinois 61801, United States; J.M. Hegstad, University of Illinois, Dept. of Crop Sciences, 1102 S. Godwin, Urbana, Illinois 618013, United States. Received 04/09/1997.

**PI 597384. Glycine max (L.) Merr.**

Cultivar. Pureline. "LN89-3615". CV-383. Pedigree - Hobbit 87 x Asgrow A3205. Indeterminate, Group IV maturity (relative maturity 4.3) maturing 6 d later than Flyer. Best adapted to 38-41 degrees N lat. Compared with Flyer averaged 5% higher yield, 45 mg seed<sup>-1</sup> larger seeds, and 10 g kg<sup>-1</sup> lower seed protein. Similar to Flyer in height, and lodging and seed quality scores. Flowers purple, pubescence tawny, pods brown and tan at maturity, and seeds shiny yellow with brown hila. Has Rps 1-k gene that confers resistance to multiple races of *P. sojae*, susceptible to brown stem rot (*Phialophora gregata*) and sudden death syndrome (*Fusarium solani*).

The following were developed by Dennis Thomas, University of Illinois, Department of Agronomy, 1102 S. Goodwin Avenue, Urbana, Illinois 61801, United States; Jay Tharp, University of Illinois, Department of Agronomy, 1102 S. Goodwin Avenue, Urbana, Illinois 61801, United States; Cecil D. Nickell, University of Illinois, Department of Crop Science, 1102 S. Goodwin Av., Urbana, Illinois 61801, United States; Greg Noel, USDA-ARS, University of Illinois, Department of Plant Pathology, Urbana, Illinois 61801, United States; T.R. Cary, University of Illinois, Illinois Agr. Exp. Sta., Dept. of Agronomy, Urbana, Illinois 61801, United States. Received 04/09/1997.

**PI 597385. Glycine max (L.) Merr.**

Cultivar. Pureline. "LN92-11008". CV-384. Pedigree - Jack x Asgrow A3205. Indeterminate, Group III maturity (relative maturity 3.1) maturing 4 d earlier than Yale and 3 d later than Jack (4,5). At SCN infested locations, seed yield was 1% higher than Yale and 2% higher than Jack. Compared with Yale at SCN noninfested locations, was similar in yield and height. Lower lodging score than Jack (1.8 vs 2.1). Flowers purple, pubescence tawny, pods tan at maturity, and seeds dull yellow with brown hila. Susceptible to *Phytophthora* rot, races 1,4, and 7 (*Phytophthora sojae*), brown stem rot (*Phialophora gregata*), and sudden death syndrome (*Fusarium solani*). Resistant to races 3 and 4 when evaluated against SCN in greenhouse.

The following were developed by Cecil D. Nickell, University of Illinois, Department of Crop Science, 1102 S. Goodwin Av., Urbana, Illinois 61801, United States. Received 04/09/1997.

**PI 597386. Glycine max (L.) Merr.**

Cultivar. Pureline. "Dwight"; LN92-10507. Pedigree - Jack x A86-303014. Indeterminate, late Group II maturity (relative maturity 2.9) maturing 5 d later than IA2021 and 1 d earlier than Jack. Compared with Jack at