

- PI 597869. *Trifolium beckwithii* W. H. Brewer ex S. Watson**
 Wild. C-103; W6 19893. Collected 08/08/1995 in California, United States . Latitude 39 deg. 59' 42'' N. Longitude 120 deg. 24' 45'' W. Elevation 1763 m. Plumas County, near town of Portola, Squaw Valley, Mercer Allotment. 9 miles from Portola (northeast). Near old railroad bed, Plumas National Forest. Dried lake bed, heavy clay, little stoniness, vernal pools/swale. Vegetation: *Artimesia arbuscula*, *Pyrrocoma hirta* spp. *lancelosa*, *Perideridia bolanterii*, *Agroseris glauca* var. *montocola*, *Iresia sericoleuca*, *Allium lemmonii*. Habit procumbent. Frequent, scattered population. *Poa secunda*, *Poa cusickii*.
- PI 597870. *Trifolium lemmonii* S. Watson**
 Wild. C-105; Lemmon's Clover; W6 19895. Collected in California, United States. Latitude 40 deg. 2' 26'' N. Longitude 120 deg. 29' 29'' W. Elevation 1794 m. Near town of Portola, Sierra Valley. South on A-23, 4-1/2 miles east off Route 70. One mile north. East exposure. 1% slope. Heavy clay (38% sand, 42% silt, 20% clay), basalt, well drained. Dry (probably wet seep in spring). Upright habit. Scattered population. Soil sample collected. Some vernal pools.
- PI 597871. *Trifolium longipes* Nutt.**
 Wild. C-106; W6 19896. Collected in California, United States. Latitude 40 deg. 2' 26'' N. Longitude 120 deg. 29' 29'' W. Elevation 1794 m. Near town of Portola, Sierra Valley. South on A-23, 4-1/2 miles east off Route 70, 1 mile north. Clay loam, basalt stones, well drained. Open exposure. Dry, probably wet seep in spring. *Perideridia golonderi* (yampa) at site. Growth upright, population frequent, scattered to dense. Flowers purple. Heads deflexed but not shattered which separates it from *lemmonii* and *beckwithii*.
- PI 597872. *Trifolium beckwithii* W. H. Brewer ex S. Watson**
 Wild. C-107; Beckwith's Clover; W6 19897. Collected in California, United States. Latitude 40 deg. 2' 26'' N. Longitude 120 deg. 29' 29'' W. Elevation 1794 m. Near town of Portola, Sierra Valley. South on A-23, 4-1/2 miles east off Route 70, 1 mile north. Clay loam, basalt stones, well drained soil. Open exposure. Dry, probably wet seep in spring. Associated species - yampa (Indian's tuber crop). East exposure, 1% slope. Upright habit. Tends to occur in swales or vernal pools whereas *T. longipes* is nearby but not in swales as much, however may co-exist in some limited areas. Heads brown, deflexed.
- PI 597873. *Trifolium wormskoldii* Lehm.**
 Wild. C-110; Wormskold's Clover; W6 19898. Collected in California, United States. Latitude 40 deg. 3' 1'' N. Longitude 120 deg. 34' 9'' W. Elevation 1702 m. Near town of Portola, Sierra Valley. South on A-23, 4-1/2 miles east off Route 70, 1 mile north. Along creek and sometimes in the creek. Associated sedges, etc. Sandy, stoney, poorly drained soil. Purple pink flower, upright growth. Occasional scattered population. Inland *T. wormskoldii* rather rare in the collection.
- PI 597874. *Trifolium longipes* Nutt.**
 Wild. C-111; Long Stalk Clover; W6 19899. Collected in California, United States. Latitude 40 deg. 2' 29'' N. Longitude 120 deg. 31' 44'' W. Elevation 1611 m. Near town of Portola, Sierra Valley. South on A-23,