

normal (9+1) di-adelphous condition. Pods green in color with dark brown streaks on surface. Seeds dark brown, round with mild shrivelling in some seeds. 100-seed mass is about 7.8g. Tolerance to wilt.

The following were developed by M.V. Reddy, Int. Crops Res. Inst. for the Semi-Arid Tropics, Crops Protection Division, Patancheru, Andhra Pradesh 502 324, India; K.B. Saxena, Int. Crops Res. Inst. for the Semi-Arid Tropics, Legumes Program, Patancheru, Andhra Pradesh 502 324, India; Laxman Singh, Int. Crops Res. Inst. for the Semi-Arid Tropics, Pigeon Pea Program, Patancheru, Andhra Pradesh 503 234, India; R.V. Kumar, Int. Crops Res. Inst. for the Semi-Arid Tropics, Patancheru P.O., Andhra Pradesh 502 324, India. Received 06/02/1997.

PI 597942. *Cajanus cajan* (L.) Millsp.

Genetic. ICPM 93003. PL-4. Pedigree - ICPM 97007/ICPL 288. Short duration indeterminate genetic male-sterile line with low fruiting branches and few secondary branches. Matures about 130 days and attains plant height of about 1.75m. Flowers yellow with red streaks on the back of standard petal and stem color green. Pods green and borne on primary branches. Produces many pods under open-pollination. Seeds cream, oval shape. Average 100-seed mass of 10.8g. Highly resistant to both wilt and sterility mosaic (SM) diseases.

The following were developed by Gary C. Peterson, Texas A&M University, Agricultural Res. & Extension Center, Route 3, Box 219, Lubbock, Texas 79401-7957, United States; Pablo Madera-Torres, USDA/ARS, Tropical Agriculture Research Station, P. O. Box 70, Mayaguez, Puerto Rico; Jeff Dahlberg, USDA, ARS, Tropical Agric. Research Station, P.O. Box 70, Mayaguez, Puerto Rico; Fred R. Miller, Texas A & M University, Department of Soil & Crop Science, College Station, Texas 77843-2474, United States; Darrell T. Rosenow, Texas Agriculture Exp. Stn., Route 3, Box 219, Farm and Market Road 1294, Lubbock, Texas 79401-9757, United States; Antonio Sotomayor-Rios, USDA, ARS, National Germplasm Repository, Tropical Agric. Research Station, Mayaguez, Puerto Rico; L.E. Clark, Texas Agr. Exp. Sta., P.O. Box 1658, Vernon, Texas 76384-1658, United States; A. Quiles-Belen, USDA, ARS, Tropical Agricultural Research Station, Box 70, Mayaguez, Puerto Rico; A. J. Hamburger, Tropical Agricultural Experiment Station, 11708 Hwy. 70 South, Vernon, Texas 76385-1658, United States; C. A. Woodfin, Tropical Agricultural Experiment Station, Route 3, Box 219, Lubbock, Texas 79401-9757, United States. Received 01/21/1997.

PI 597943. *Sorghum bicolor* (L.) Moench

Breeding. Pureline. IS 1117C; PI 248313 (preconverted); SC 196; Satsemari. GP-520. Pedigree - Photoperiod sensitive conversion of IS 1117 selected from the World Sorghum Collection. Early maturing, combine-height enhanced. Tested for fertility reactions in A1 cytoplasm and found to be a partial maintainer line. A unique Durra (race) - Durra (Working Group 41).

PI 597944. *Sorghum bicolor* (L.) Moench

Breeding. Pureline. IS 2680C; PI 267366 (preconverted); SC 808; J-16. GP-521. Pedigree - Photoperiod sensitive conversion of IS 2680 selected from the World Sorghum Collection. Early maturing, combine-height