

- PI 598028. *Glycine max* (L.) Merr.**
Genetic. Pureline. A96-1232-3. GP-235. Pedigree - ms6 ms6 w1 w1 x AX2858 . Near-isogenic line cosegregating M6 W1 ms6 w1. The Ms6 and W1 alleles in coupling phase are tightly linked (2 to 5% genetic recombination). Plants self-pollinated to allow maintenance of the two alleles. Plants identical to AX2858 in all phenotypic traits measured except for the Ms6 and W1 loci.
- PI 598029. *Glycine max* (L.) Merr.**
Genetic. Pureline. A96-1283-3. GP-236. Pedigree - ms6 ms6 w1 w1 x A3307. Near-isogenic line Ms6 ms6 w1 w1. Plants self-pollinated to identify segregation of the ms6 allele. Plants identical to A3307 in all phenotypic traits measured except for the Ms6 locus.
- PI 598030. *Glycine max* (L.) Merr.**
Genetic. Pureline. A96-1298-4. GP-237. Pedigree - ms6 ms6 w1 w1 x A3307. Near-isogenic line Ms6 ms6 w1 w1. Plants self-pollinated to identify segregation of the ms6 allele. Plants identical to A3307 in all phenotypic traits measured except for the Ms6 locus.
- PI 598031. *Glycine max* (L.) Merr.**
Genetic. Pureline. A96-989-2. GP-238. Pedigree - ms6 ms6 w1 w1 x CX-155. Near-isogenic line cosegregating M6 W1 ms6 w1. The Ms6 and W1 alleles in coupling phase are tightly linked (2 to 5% genetic recombination). Plants self-pollinated to allow maintenance of the two alleles. Plants identical to CX-155 in all phenotypic traits measured except for the Ms6 and W1 loci.
- PI 598032. *Glycine max* (L.) Merr.**
Genetic. Pureline. A96-1154-4. GP-239. Pedigree - ms6 ms6 w1 w1 x CX-155 . Near-isogenic line cosegregating M6 W1 ms6 w1. The Ms6 and W1 alleles in coupling phase are tightly linked (2 to 5% genetic recombination). Plants self-pollinated to allow maintenance of the two alleles. Plants identical to CX-155 in all phenotypic traits measured except for the Ms6 and W1 loci.
- PI 598033. *Glycine max* (L.) Merr.**
Genetic. Pureline. A96-1314-2. GP-240. Pedigree - ms6 ms6 w1 w1 x 82-165 . Near-isogenic line cosegregating M6 W1 ms6 w1. The Ms6 and W1 alleles in coupling phase are tightly linked (2 to 5% genetic recombination). Plants self-pollinated to allow maintenance of the two alleles. Plants identical to 82-165 in all phenotypic traits measured except for the Ms6 and W1 loci.
- PI 598034. *Glycine max* (L.) Merr.**
Genetic. Pureline. A96-1335-4. GP-241. Pedigree - ms6 ms6 w1 w1 x 82-165 . Near-isogenic line cosegregating M6 W1 ms6 w1. The Ms6 and W1 alleles in coupling phase are tightly linked (2 to 5% genetic recombination). Plants self-pollinated to allow maintenance of the two alleles. Plants identical to 82-165 in all phenotypic traits measured except for the Ms6 and W1 loci.
- PI 598035. *Glycine max* (L.) Merr.**
Genetic. Pureline. A96-1346-2. GP-242. Pedigree - ms6 ms6 w1 w1 x 82-378