

The following were collected by David E. Williams, USDA, ARS, Natl. Germplasm Resources Laboratory, Building 003, Room 400, BARC-West, Beltsville, Maryland 20705-2350, United States. Donated by Thomas G. Isleib, North Carolina State University, Department of Crop Science, Box 5155, Raleigh, North Carolina 27695-7629, United States; David E. Williams, USDA, ARS, Natl. Germplasm Resources Laboratory, Building 003, Room 400, BARC-West, Beltsville, Maryland 20705-2350, United States. Received 07/14/1997.

PI 599606. *Arachis hypogaea* var. *fastigiata* (Waldron) Krapov. & W. C. Greg.

Landrace. DEW 1010; mani de cascara negra. Collected 09/25/1989 in Beni, Bolivia. Latitude 14 deg. 30' 0'' S. Longitude 67 deg. 30' 0'' W. Elevation 320 m. Dept. Beni, Prov. Ballivian, Carmen Florida, 7 km upriver from Rurrenabaque on right bank of the Rio Beni. Small village of acculturated Tacana and Quechua natives. Cultivated in chaco (swidden field) in upland forest. Plants erect. Flowers orange. Fruits large, turning dark brown or black when mature, containing 4 red seeds. Pegs purple. Once-common local landrace now rare.

PI 599607. *Arachis hypogaea* var. *fastigiata* (Waldron) Krapov. & W. C. Greg.

Landrace. DEW 925; mani colorado. Collected 06/10/1989 in Santa Cruz, Bolivia. Latitude 17 deg. 45' 0'' S. Longitude 63 deg. 22' 0'' W. Elevation 400 m. Dept. Santa Cruz, Prov. Andres Ibanez, Canton Terebinto, 35km W of Santa Cruz. Cultivated on Lic. G. Coimbra Sanz's property. Pods fairly straight, fairly smooth with slight constriction, small humps, slight beak, containing 3 red seeds. Local variety traditionally grown in the area.

PI 599608. *Arachis hypogaea* var. *fastigiata* (Waldron) Krapov. & W. C. Greg.

Landrace. DEW 928; mani. Collected 06/12/1989 in Beni, Bolivia. Latitude 14 deg. 30' 0'' S. Longitude 67 deg. 30' 0'' W. Elevation 268 m. Dept. Beni, Prov. Ballivian, Rurrenabaque. Purchased at corner store, Calle Comercio. Cultivated locally. Mix of two kinds of fruits. Some reticulated and some almost or completely devoid of reticulation. Reticulated pods fairly straight, with humps, some beak, little or no constriction, containing 3-4 predominantly red, some white seeds. Non-reticulated pods very smooth, sometimes enlarged at base, with humps, beak, little or no constriction, containing 3-4 almost exclusively red seeds, with a very few pods containing white seeds. Fruits dusty, suggesting chaco rather than sandy playa cultivation. White seeds now turned tan with age. Probably from last year's harvest.

PI 599609. *Arachis hypogaea* var. *fastigiata* (Waldron) Krapov. & W. C. Greg.

Landrace. DEW 930; mani negro. Collected 07/19/1989 in Beni, Bolivia. Latitude 14 deg. 30' 0'' S. Longitude 67 deg. 30' 0'' W. Elevation 268 m. Dept. Beni, Prov. Ballivian, Rurrenabaque. Cultivated locally. Fruit straight, with humps, beak, little or no constriction, containing 3-4 very dark purple seeds. Only 3 such fruits were found when shelling part of sample #929.