

32917 to 32973—Continued.

- 32921.** BERBERIS GUIMPELI Koch and Bouche. **Barberry.**
- 32922.** BERBERIS JAPONICA BEALEI (Fortune) Skeels. **Barberry.**
See Nos. 31244 and 32700 for previous introductions.
- 32923.** BERBERIS NAPAULENSIS (DC.) Spreng. **Barberry.**
See No. 28839 for description.
- 32924.** CAJUPUTI CUTICULARIS (Labill.) Skeels.
(*Melaleuca cuticularis* Labill. 1806, *Novae Hollandiae Plantarum Specimen*, vol. 2, p. 30, pl. 171.)
- 32925.** CAJUPUTI ERICIFOLIA (Smith) Lyons. **Bottle-brush tea-tree.**
See No. 30793 for previous introduction.
- 32926.** CAJUPUTI HYPERICIFOLIA (Salisb.) Skeels. **Hillock tree.**
See No. 30761 for previous introduction.
- 32927.** CAJUPUTI PUBESCENS (Schauer) Skeels.
See No. 30795 for previous introduction.
- 32928.** CAJUPUTI PULCHELLA (R. Br.) Skeels.
(*Melaleuca pulchella* R. Br. 1812, *Aiton Hortus Kewensis*, ed. 2, vol. 4, p. 414.)
- 32929.** CAJUPUTI WILSONI (Mueller) Skeels. **Wilson's tea-tree.**
(*Melaleuca wilsoni* Mueller 1861, *Fragmenta Phytographiae Australiae*, vol. 2, p. 124, pl. 15.)

The reason for the use of the generic name *Cajuputi* in place of *Melaleuca* is explained in Inventory No. 27 (*Bulletin* 242, Bureau of Plant Industry, U. S. Dept. of Agriculture, 1912, p. 39.) *Cajuputi cuticularis* is a tall shrub or small tree and *C. pulchella* is a low, spreading shrub, both growing along streams in West Australia. *C. wilsoni* is a tall shrub found in Victoria and South Australia.

- 32930.** CELTIS AUDIBERTIANA Spach. **Hackberry.**
This species is generally referred to *C. occidentalis* L. As the latter is now considered to cover several distinct forms it is thought best to retain the present name until the material can be grown.
- 32931.** COLUTEA CILICICA Boiss. and Bal.
Distribution.—A shrub found on the coast and on the slopes of the mountains in the southern part of Asia Minor from Smyrna to Cilicia.
- 32932.** COTONEASTER AFFINIS Lindl.
See No. 28207 for previous introduction.
- 32933.** COTONEASTER BUXIFOLIA Wall.
Distribution.—On the slopes of the Nilgiri Hills in the southern part of India.
- 32934.** COTONEASTER MICROPHYLLA Wall.
Distribution.—A shrub found on the temperate slopes of the Himalayas at an altitude of 4,000 to 8,000 feet, between Kashmir and Bhutan in northern India.
- 32935.** COTONEASTER MICROPHYLLA THYMIFOLIA (Loud.) Koehne.
Distribution.—A low shrub found on the alpine slopes of the Himalayas from Kashmir to Sikkim in northern India.
- 32936.** COTONEASTER PANNOSA Franchet.
Distribution.—A low shrub found on limestone ledges of the mountains in the province of Yunnan in China at an elevation of 10,000 feet.