

38548 to 38567—Continued. (Quoted notes by Mr. O. F. Cook.)

38566. *ACHRADELPHA VIRIDIS* (Pittier) O. F. Cook. Sapotaceæ.
(*Calocarpum viride* Pittier.)

“Inferior variety from the market of the city of Guatemala.”

See S. P. I. Nos. 38478 to 38481 for previous introductions and description.

38567. *CHAYOTA EDULIS* Jacq. Cucurbitaceæ. **Chayote.**
(*Sechium edule* Sw.)

“Thin-skinned variety of chayote from the market of the city of Guatemala.”

38568. *ERIOBOTRYA JAPONICA* (Thunb.) Lindl. Malaceæ. **Loquat.**

From Algiers, Algeria. Presented by Dr. L. Trabut. Received June 20, 1914.

“*Tanaka* loquat. An excellent variety of fruit, with very firm flesh of a yellow color. Stands transportation for a period of one week. In Algeria the seeds give varieties superior to the original type, furnishing interesting varieties for the market.” (*Trabut.*)

See S. P. I. No. 8890 for description of the original introduction of this variety from Japan into the United States.

38569 and 38570. *HOLCUS SORGHUM* L. Poaceæ. **Sorghum.**
(*Sorghum vulgare* Pers.)

From Victoria, Kamerun, German West Africa. Presented by the director of the Experiment Station. Received June 1, 1914. Quoted notes by the director.

“Seed from the Mora residency in the German lands near Lake Chad.”

38569. “*Gabli nialgo*. Sown at the rainy season.”

38570. “*Massaggioa adjagama*. For the manufacture of firkiboden this variety is preferred. It is planted after the rainy season.”

38571 to 38576. *RUBUS* spp. Rosaceæ.

From Sibpur, near Calcutta, India. Presented by the Royal Botanic Gardens. Received June 15, 1914.

38571. *RUBUS ALPESTRIS* Blume.

Distribution.—A straggling shrub found on the temperate slopes of the Himalayas at an altitude of 7,000 to 18,000 feet and also in Java.

38572. *RUBUS ANDERSONI* Hook. f.

Distribution.—A bramble from altitudes of 7,000 to 8,000 feet in the Sikkim Himalayas.

38573. *RUBUS CALYCNUS* Wallich.

Distribution.—A creeping herbaceous perennial with simple reniform leaves and scarlet fruits, usually with but few fruitlets, found in India on the temperate slopes of the Khasi Hills at an altitude of 4,000 to 5,000 feet.

38574. *RUBUS NIVEUS* Thunb.

See S. P. I. Nos. 32453, 33344, and 34334 for previous introductions and description under the name *R. lasiocarpus*.