

180912 to 180949—Continued

- 180934. *Lischower Früh.*
- 180935. *Lohmanns Weender IV.*
- 180936. *Maurener Weibhafer V. Reininghaus.*
- 180937. *Norddeutsche Pflanzenzucht (NPZ).* Seeds white.
- 180938. *Peragis Früh II.*
- 180939. *Peragis Weib.*
- 180940. *Pfetten Weihenstephaner.*
- 180941. *PSG Goldkorn (Pommersche Saatzuchtgesellschaft).*
- 180942. *Rotenburger Schwarz.*
- 180943. *Siegeshafer.*
- 180944. *Streckenthiner 9.*
- 180945. *Strubes Gelb II.*
- 180946. *Svalöfs Adler.*
- 180947. *Svalöfs Goldregen III.*
- 180948. *Wadsacks Gelb.*
- 180949. *Wentzels Anderbecker Gelb.*

180950 to 180952. GOSSYPIUM HERBACEUM L. Malvaceae. Cotton.

From India. Seeds presented by the Office of the Cotton Botanist, Surat. Received Apr. 21, 1949.

- 180950. No. 1027. A. L. F.
- 180951. 1A *Long Boll.*
- 180952. *Vijaya.*

180953 to 180955. FRAGARIA. Rosaceae. Strawberry.

From Portugal. Plants presented by Alfredo Moreira da Silva & Filhos, Porto. Received Apr. 21, 1949.

- 180953. *Santo Tirso.*
- 180954. *Porto.*
- 180955. *Matosinkos.*

180956 to 181007.

From India. Seeds collected by Walter N. Koelz, Agricultural Explorer, Bureau of Plant Industry, Soils, and Agricultural Engineering, Beltsville, Md. Received Apr. 21, 1949.

Collected in 1948 unless otherwise indicated.

180956 to 181001. TRITICUM AESTIVUM L. Poaceae. Common wheat.

- 180956 to 180965. From the region of the eastern Punjab near the Tibetan border.
- 180956. 8-A.
- 180957. 9-D. Early type.
- 180958. C-217.
- 180959. C-228. Adapted to late sowing.
- 180960. C-250. Rust-resistant.
- 180961. C-259.
- 180962. C-260.
- 180963. C-275.
- 180964. C-518. Strong stems; yields reported to be 4,480 pounds per acre.
- 180965. C-591. Major wheat type of Punjab Province.