

296988 to 297000. HORDEUM VULGARE L. Gramineae.

From Argentina. Seeds presented by the Instituto Nacional de Tecnología Agropecuaria, Bordenave, via the University of Oklahoma, Stillwater, Okla. Received Apr. 7, 1964.

Varieties screened for greenbug resistance rated on damage as follows: 3=41-60 percent, 4=61-80 percent, 5=beyond recovery.

<i>P.I. No.</i>	<i>Strain</i>	<i>Source</i>	<i>Rating</i>	<i>Number of living plants</i>
296988.	B.216-550100b	2884/62	5	1
296989.	B.216-6202	2889/62	5	2
296990.	B.217-1503	2912/62	5	1
296991.	B.217-2302	2913/62	5	3
296992.	B.217-4201	2914/62	5	1
296993.	B.217-7501	2922/62	5	2
296994.	B.217-(11)40(12)	2925/62	5	2
296995.	B.273-0(15)000	3245/62	4	5
296996.	B.273-0(15)001	3246/62	4	7
296997.	B.273-0(15)002	3247/62	3	15
296998.	B.273-0(26)00a3	3286/62	5	2
296999.	B.273-0(26)00a5	3294/62	5	2
297000.	B.273-0(29)000	3302/62	5	1

297001 to 297030.

From Kenya. Seeds presented by the Plant Breeding Station, Njoro. Received Apr. 7, 1964.

297001. TRITICUM (hybrid). Gramineae.

No. 8. 'Mengavi' ('Spica-Koda' × 'Gabo'). 'Gabo' type resistance. Pedigree: 'Spica', 'Gabo', and *T. timopheevi*.

297002. TRITICUM (hybrid).

No. 20. *T. aestivum* × *Aegilops ovata*. *Aegilops* resistance.

297003 to 297029. TRITICUM AESTIVUM L. Common wheat.

297003. 'Yaqui 50'. 'Marroqui 588' × 'Newthatch'.

297004. 'Kentana Yaqui'.

297005. 'Africa-Mayo 48'.

297006. 1066/1. Unknown, field selection.

297007. 'MidalmcMurachy' × 'Exchange'.

297008. 'Veranopolis'. Unknown. Susceptible to race 34 in Kenya.

297009. 'Tama'. 'Yaktana' × 'Lerma'.

297010. 'Gala'. Spring wheat—'Gabo' × 'Lerma'. 'Gabo' type resistance.

297011. 'Yaqui 53'. Y.50 × Eg.101 × 'Gabo'. 'Gabo' type resistance.

297012. 'Yansek' × 'Bonza'. 'Bonza' type resistance.

297013. T-K² × 'Bonza² VI.106'. 'Bonza' type resistance.

297014. 'Fronthatch'. 'Frontana-K.58-Newthatch'. 'Thatcher' type resistance.

297015. 'Morris'. (Mida-Kenya 117A-Thatcher²-Frontana-Thatcher⁴). 'Thatcher' type resistance.