

343231 to 343233—Continued

- 343231.** Jutiapa race. Edge of field near El Progreso South, Jutiapa. Elevation 1,100 m. Plant 144 cm. high, 9.3 tillers, 11.4 nodes, green, sheaths glabrous; 90 days to flower. January 1964.
- 343232.** Huehuetenango race. Canyon near San Antonio Hiusta, Huehuetenango. Elevation 1,300 m. Plant 173 cm. high, 11.5 tillers, 14.3 nodes, green to reddish, slight pubescence; 104 days to flower. January 1964.
- 343233.** Huehuetenango race. Along trail at Tzisbaj, Huehuetenango. Elevation 1,510 m. Plant 183 cm. high, 13 tillers, 15.3 nodes, green to reddish, slight pubescence; 104 days to flower. February 1964.

343234 to 343249. EUCHLAENA MEXICANA Schrad. Gramineae.
Teosinte.

From Mexico. Seed collected by H. G. Wilkes, Department of Biology, Tulane University, New Orleans, Louisiana. Numbered June 3, 1969.

- 343234.** 45121. Central Plateau race. Edge of field near Cerro 8 km. east of Uriangato, Guanajuato. Elevation 1,850 m. Plant 139 cm. high, 3.9 tillers, 11.4 nodes, green to reddish, slightly pubescent; 78 days to flower. November 1963.
- 343235.** 45320. Central Plateau race. Edge of field, Churuntzio, Michoacan. Elevation 1,900 m. Plant 148 cm. high, 7.7 tillers, 13.3 nodes, green to reddish, slightly pubescent; 79 days to flower. November 1963.
- 343236.** 46452. Central Plateau race. Edge of field, Manual Doblado, Guanajuato. Elevation 1,750 m. Plant 169 m. high, 4.9 tillers, 12.9 nodes, green to reddish, slightly pubescent; 79 days to flower. December 1963.
- 343237.** 47259. Balsas race. Mountainside, Palo Blanco, Guerrero. Elevation 1,250 m. Plant 114 cm. high, 11.6 tillers, 11.9 nodes, green to reddish, sheath nearly glabrous; 91 days to flower. December 1963.
- 343238.** 47269. Balsas race. Limestone ledge, Acshuizotla, Guerrero. Elevation 1,100 m. Plant 110 cm. high, 13.8 tillers, 11.8 nodes, green to reddish, sheath nearly glabrous; 87 days to flower. December 1963.
- 343239.** 47335. Balsas race. Pasture area, Mazatlan, Guerrero. Elevation 1,350 m. Plant 106 cm. high, 12.5 tillers, 11.1 nodes, green to reddish, sheath nearly glabrous; 90 days to flower. December 1963.
- 343240.** 47711. Balsas race. Mountainside near fields, Cerro Ranchos Nuevos on road Teloloapan-Arcelia, Guerrero. Elevation 1,510–1,900 m. Plant 165 cm. high, 6.6 tillers, 12.9 nodes, green to reddish, sheath nearly glabrous; 86 days to flower. December 1963.
- 343241.** 47813. Balsas race. Mountain side, Cerro de Los Chivos on road Teloloapan-Arcelia, Guerrero. Elevation 840 m. Plant 120 cm. high, 9.5 tillers, 11.3 nodes, green to reddish, sheath nearly glabrous; 88 days to flower. December 1963.
- 343242.** 47890. Balsas race. Cerro east of Huetamo, Michoacan. Elevation 600–800 m. Plant 25 cm. high, 8.4 tillers, 10.9 nodes, green to reddish, sheath nearly glabrous; 92 days to flower. December 1963.
- 343243.** 48085. Balsas race. Queretanillo on bank of Rio Cutzamala on Huetamo-Morelia road, Michoacan. Elevation 800 m. Plant 123 cm. high, 8.7 tillers, 12.2 nodes, green to reddish, sheath nearly glabrous; 86 days to flower. December 1963.
- 343244.** 48703. Central Plateau race. Edge of field, Copandirrom, Michoacan. Elevation 1,850 m. Plant 170 cm. high, 6.0 tillers, 14 nodes, green to reddish, slightly pubescent; 79 days to flower. December 1963.
- 343245.** 49094. Balsas race. Cornfield near Taxco, Guerrero. Elevation 2,350–2,500 m. Plant 171 cm. high, 5.4 tillers, 13.6 nodes, green to reddish, sheath nearly glabrous; 89 days to flower. December 1963.