

PI 511897 to 511936-continued

- PI 511913 **donor id:** 851124-04. **origin:** Mexico. **local name:** Arota. **collected:** November 24, 1985. **collector id:** 851124-04. **locality:** Level site on hilly farmland of Rancho del Babo, 39km NW of El Fuerte, Rio Fuerte region, Sinaloa. **latitude:** 26 deg. 35 min. N. **longitude:** 108 deg. 20 min. W. **elevation:** 320m. **remarks:** Plant more adaptable to varying soil types than Cucurbita moschata. Sown in June and July, harvested in October and November. Cook mature fruits. Sample donated by Encornacion Torres Legarra. Primitive Cultivar. Seed.
- PI 511914 **donor id:** 851124-06. **origin:** Mexico. **local name:** Arota. **collected:** November 24, 1985. **collector id:** 851124-06. **locality:** Level site on hilly farmland of Rancho del Babo, 39km NW of El Fuerte, Rio Fuerte region, Sinaloa. **latitude:** 26 deg. 35 min. N. **longitude:** 108 deg. 20 min. W. **elevation:** 320m. **remarks:** Plant more adaptable to varying soil types than Cucurbita moschata. Sown in June and July, harvested in October and November. Cook mature fruits. Sample donated by Encornacion Torres Legarra. Primitive Cultivar. Seed.
- PI 511915 **donor id:** 851124-11. **origin:** Mexico. **local name:** Arota (con cascara dura). **collected:** November 24, 1985. **collector id:** 851124-11. **locality:** Hilly farmland of Rancho Guayobitas, 9km N of Choix, Rio Fuerte region, Sinaloa. **latitude:** 26 deg. 45 min. N. **longitude:** 108 deg. 30 min. W. **elevation:** 260m. **remarks:** Plant may be a cross between Cucurbita mixta and Cucurbita palmeri due to presence of hard rind. Sown in July, harvested in October and November. Cook mature flesh as a sweet, toast or grind seeds for a sauce (Pipian). Sample donated by Manuel Pina Corrales. Primitive Cultivar. Seed.
- PI 511916 **donor id:** 851124-12. **origin:** Mexico. **local name:** Arota. **collected:** November 24, 1985. **collector id:** 851124-12. **locality:** Hilly farmland of Rancho Guayobitas, 9km N of Choix, Rio Fuerte region, Sinaloa. **latitude:** 26 deg. 45 min. N. **longitude:** 108 deg. 30 min. W. **elevation:** 260m. **remarks:** Fruits diverse in shape and color. Sown in July, harvested in October and November. Cook mature flesh as a sweet, toast or grind seeds for a sauce (Pipian). Sample donated by Manuel Pina Corrales. Primitive Cultivar. Seed.